

## **A functional upper extremity after stroke: scientific evidence applied in the current health care climate**

**Patricia S. Pohl, PhD, PT**  
**Department of Physical Therapy & Rehabilitation Sciences**  
**University of Kansas Medical Center**

### **References**

An-Yeung, S. S., Ng, J. T., & Lo, S. K. (2003). Does balance or motor impairment of limbs discriminate the ambulatory status of stroke survivors. American Journal of Physical Medicine and Rehabilitation, *82*, 279-283.

Bonifer, N., & Anderson, K. M. (2003). Application of constraint-induced movement therapy for an individual with severe chronic upper-extremity hemiplegia. Physical Therapy, *83*, 384-398.

Butefisch, C., Hummelsheim, H., Denzler, P., & Mauritz, K.-H. (1995). Repetitive training of isolated movements improves the outcome of motor rehabilitation of the centrally paretic hand. Journal of the Neurological Sciences, *130*, 59-68.

Cambier, D. C., De Corte, E., Danneels, L. A., & Witvrouw, E. E. (2003). Treating sensory impairments in the post-stroke upper limb with intermittent pneumatic compression. Results of a preliminary trial. Clinical Rehabilitation, *17*, 14-20.

Cauraugh, J. H., & Kim, S. (2003). Progress toward motor recovery with active neuromuscular stimulation: muscle activation pattern evidence after a stroke. Journal of Neurological Science, *207*, 25-29.

Cunningham, C. L., Stoykov, M.E., & Walter, C. B. (2002). Bilateral facilitation of motor control in chronic hemiplegia. Acta Psychologica, *110*, 321-337.

de Kroon, J. R., van der Lee, J. H., Ijzerman, M. J., & Lankhorst, G. J. (2002). Therapeutic electrical stimulation to improve motor control and functional abilities of the upper extremity after stroke: a systematic review. Clinical Rehabilitation, *16*, 350-360.

Desrosiers, J., Malouin, F., Richards, C., Bourbonnais, D., Rochette, A., & Bravo, G. (2003). Comparison of changes in upper and lower extremity impairments and disabilities after stroke. International Journal of Rehabilitative Research, *26*, 109-116.

Duncan, P., Studenski, S., Richards, L., Gollub, S., Lai, S. M., Reker, D. et al. (2003). Randomized clinical trial of therapeutic exercise in subacute stroke. Stroke, *34*, 2173-2180.

Fasoli, S. E., Krebs, H. I., Stein, J., Frontera, W. R., & Hogan, N. (2003). Effects of robotic therapy on motor impairment and recover in chronic stroke. Archives of Physical Medicine and Rehabilitation, 84, 477-482.

Hesse, S., Schulte-Tiggles, G., Konrad, M., Bardeleben, A., & Werner, C. (2003). Robot-assisted arm trainer for the passive and active practice of bilateral forearm and wrist movements in hemiparetic patients. Archives of Physical Medicine and Rehabilitation, 84, 915-920.

Kemper, S., McDowd, J., Pohl, P., Herman, R., & Jackson, S. (in press). Revealing language deficits following stroke: the cost of doing two things at once. Aging, Neuropsychology and Cognition.

Kozlowski, D. A., James, D. C., & Schallert, T. (1996). Use-dependent exaggeration of neuronal injury after unilateral sensorimotor cortex lesions. The Journal of Neuroscience, 16, 4776-4786.

Lai, S. M., Studenski, S., Duncan, P. W., & Perera, S. (2002). Persisting consequences of stroke measured by the Stroke Impact Scale. Stroke, 33, 1840-1844.

Liepert, J., Terborg, C., & Weiller, C. (1999). Motor plasticity induced by synchronized thumb and foot movements. Experimental Brain Research, 125, 435-439.

Lum, P. S., Burgar, C. G., Shor, P. C., Majmundar, M., & Van der Loos, M. (2002). Robot-assisted movement training compared with conventional therapy techniques for the rehabilitation of upper-limb motor function after stroke. Archives of Physical Medicine and Rehabilitation, 83, 952-959.

Ma, H., Trombly, C. A., & Robinson-Podolski, C. (1999). The effect of context on skill acquisition and transfer. American Journal of Occupational Therapy, 53, 138-144.

Merians, A. S., Jack, D., Boian, R., Tremaine, M., Burdea, G. C., Adamovich, S. V., Recce, M., & Poizner, H. (2002). Virtual reality-augmented rehabilitation for patients following stroke. Physical Therapy, 82, 898-915.

Mudie, M. H., & Matyas, T. A. (2001). Responses of the densely hemiplegic upper extremity to bilateral training. Neurorehabilitation and Neural Repair, 15, 129-140.

Nudo, R. J., Wise, B. M., SiFuentes, F., & Milliken, G. W. (1996). Neural substrates for the effects of rehabilitation training on motor recovery after ischemic infarct. Science, 272, 1791-1794.

Page, S. J., Levine, P., Sisto, S., Bond, Q., & Johnston, M. V. (2002). Stroke patients' and therapists' opinions of constraint-induced movement therapy. Clinical Rehabilitation, 16, 55-60.

Page, S. J., Sisto, S. A., & Levine, P. (2002). Modified constraint-induced therapy in chronic stroke. American Journal of Physical Medicine and Rehabilitation, *81*, 870-875.

Pandyan, A. D., Cameron, M., Powell, J., Stott, D. J., & Granat, M. H. (2003). Contractures in the post-stroke wrist: a pilot study of its time course of development and its association with upper limb recovery. Clinical Rehabilitation, *17*, 88-95.

Pohl, P. S., Winstein, C. J., & Onla-Or, S. (1997). Sensory-motor control in the ipsilesional upper extremity after stroke. NeuroRehabilitation, *9*, 57-69.

Pons, T. P., Garraghty, P. E., Ommaya, K., Kaas, J. H., Taub, E., & Mishkin, M. (1991). Massive cortical reorganization after sensory deafferentation in adult macaques. Science, *252*, 1857-1860.

Priplata, A., Niemi, J., Salen, M., Harry, J., Lipsitz, L. A., & Collins, J. J. (2002). Noise-enhanced human balance control. Physics Review Letters, *89*, 238101.

Reinkensmeyer, D. J., Pang, C. T., Nessler, J. A., & Painter, C. C. (2002). Web-based telerehabilitation for the upper extremity after stroke. IEEE Trans Neural Systems Rehabilitation Engineering, *10*, 102-108.

Riolo, L., & Fisher, K. (2003). Evidence in practice. Physical Therapy, *83*, 844-851.

Steenbergen, B., Hulstijn, W., de Vries, A., & Berger, M. (1996). Bimanual movement coordination in spastic hemiparesis. Experimental Brain Research, *110*, 91-98.

Stelmach, G. E. (1969). Efficiency of motor learning as a function of intertrial rest. Research Quarterly, *40*, 198-202.

Stevens, J. A., & Stoykov, M. E. (2003). Using motor imagery in the rehabilitation of hemiparesis. Archives of Physical Medicine and Rehabilitation, *84*, 1090-1092.

Taub, E., Crago, J. E., Burgio, L. D., Groomes, T. E., Cook III, E. W., DeLuca, S. C., & Miller, N. E. (1994). An operant approach to rehabilitation medicine: overcoming learned nonuse by shaping. Journal of the Experimental Analysis of Behavior, *61*, 281-293.

Taub, E., & Uswatte, G. (2003). Constraint-induced movement therapy: bridging from the primate laboratory to the stroke rehabilitation laboratory. Journal of Rehabilitation Medicine, *Suppl. 41*, 34-40.

Tillerson, J. L., Cohen, A. D., Caudle, W. M., Zigmond, M. J., Schallert, T., & Miller, G. W. (2002). Forced nonuse in unilateral Parkinsonian rats exacerbates injury. The Journal of Neuroscience, *22*, 6790-6799.

Van der Lee, J. H. (2003). Constraint-induced movement therapy: some thoughts about theories and evidence. Journal of Rehabilitation Medicine, *Supplement 41*, 41-45.

Volpe, B. T., Krebs, H. I., & Hogan, N. (2003). Robot-aided sensorimotor training in stroke rehabilitation. Advances in Neurology, *92*, 429-433.

West, S. K., Rubin, G. S., Munoz, B., Abraham, D., Fried, L. P., & Salisbury Eye Evaluation Project Team (1997). Assessing functional status: correlation between performance on tasks conducted in a clinic setting and performance on the same task conducted at home. Journal of Gerontology: Medical Science, *52A*, M209-M217.

Whitall, J., Waller, S. M., Silver, K. H. C., & Macko, R. F. (2000). Repetitive bilateral arm training with rhythmic auditory cueing improves motor function in chronic hemiparetic stroke. Stroke, *31*, 2390-2395.

Woldag, H., & Hummelsheim, H. (2002). Evidence-based physiotherapeutic concepts for improving arm and hand function in stroke patients. A review. Journal of Neurology, *249*, 518-528.

Wolf, S. L., Lecraw, E., Barton, L. A., & Jann, B. B. (1989). Forced use of hemiplegic upper extremities to reverse the effect of learned nonuse among chronic stroke and head-injured patients. Experimental Neurology, *104*, 125-132.

Wright, D. L., & Shea, C. H. (1991). Contextual dependencies in motor skills. Memory & Cognition, *19*, 361-370.

Wu, C., Trombly, C. A., Lin, K., & Tickle-Degnen, L. (2000). A kinematic study of contextual effects on reaching performance in persons with and without stroke: influences of object availability. Archives of Physical Medicine and Rehabilitation, *81*, 95-101.